

SUPPLEMENTARY DATA

Supplementary Table 1. Baseline Characteristics by Frequency of Statin use During the Study

	≤ 50% of visits (n=36)	> 50% of visits (n=161)
Age (years)	62 ± 11	61 ± 9
Diabetes Duration (years)	11 ± 8	12 ± 8
Non Hispanic Whites vs. Others (%)	36	32
Hypertension (%)	75	82
Prior CVD (%)	19	41*
BMI (kg/m ²)	32.2 ± 4.4	31.1 ± 4.5
Systolic Blood Pressure	130 ± 15	132 ± 17
Diastolic Blood Pressure	74 ± 9	76 ± 10
HbA1C (%)	9.4 ± 1.7	9.1 ± 1.2
Total Cholesterol (mg/dl)	179 ± 47	181 ± 38
LDL-C (mg/dl)	110 ± 29	104 ± 30
HDL-C (mg/dl)	37 ± 10	37 ± 11
Triglycerides (mg/dl)	183 (124 -235)	159 (123 -236)
Total-C/HDL	5.1 ± 1.3	5.2 ± 1.7
Triglycerides/HDL	6.6 ± 5.4	6.2 ± 5.4
CAC (mm ³)	16.1 ± 17.8	18.5 ± 15.3
AAC (mm ³)	32.7 ± 33.5	37.1 ± 29.2
Intensive treatment	47	47

Number are mean (± SD) or median(25th-75th percentiles) or frequencies; Volume square root transformed values for CAC and AAC are presented CAC and AAC

* $P < 0.01$