

SUPPLEMENTARY DATA

Supplementary Table 1. Percentage of the population by risk score category and age category *

| Score category | Age category (years) | | | | | | | |
|----------------|----------------------|---------|---------|---------|---------|---------|---------|---------|
| | < 45 | 45-49.9 | 50-54.9 | 55-59.9 | 60-64.9 | 65-69.9 | 70-74.9 | 75-85.9 |
| Males | | | | | | | | |
| < 25 | 98.4 | 72.2 | 47.7 | 15.5 | - | - | - | - |
| 25 – 29 † | 1.6 | 18.2 | 25.2 | 29.3 | - | - | - | - |
| 30 – 34 | - | 5.1 | 19.8 | 21.7 | 12.2 | - | - | - |
| 35 – 39 | - | 2.8 | 4.5 | 23.0 | 20.2 | 14.2 | - | - |
| 40 – 44 | - | 0.6 | 2.7 | 4.4 | 28.6 | 25.5 | - | - |
| 45 – 49 | - | 1.1 | - | 5.3 | 20.6 | 21.4 | 15.8 | - |
| 50 - 54 | - | - | - | 0.7 | 10.3 | 25.3 | 32.6 | - |
| 55 - 59 | - | - | - | 0.2 | 5.1 | 7.7 | 19.7 | - |
| ≥ 60 | - | - | - | - | 2.9 | 5.8 | 31.9 | 100 |
| Females | | | | | | | | |
| < 25 | 97.6 | 75.6 | 40.6 | 8.7 | - | - | - | - |
| 25 – 29 † | 1.8 | 13.0 | 30.9 | 23.7 | 8.0 | - | - | - |
| 30 – 34 | 0.4 | 7.3 | 13.3 | 22.8 | 11.7 | - | - | - |
| 35 – 39 | 0.2 | 2.6 | 6.1 | 19.7 | 26.5 | 10.5 | - | - |
| 40 – 44 | - | 1.6 | 7.3 | 13.8 | 20.5 | 17.0 | - | - |
| 45 – 49 | - | - | 1.8 | 7.8 | 13.3 | 20.4 | 5.0 | - |
| 50 - 54 | - | - | - | 2.3 | 13.9 | 23.6 | 9.5 | - |
| 55 - 59 | - | - | - | 1.2 | 4.1 | 12.6 | 28.2 | - |
| ≥ 60 | - | - | - | - | 2.0 | 15.9 | 57.4 | 100 |

* Data from the Rotterdam Study and PREVEND were used for this Table. Hoorn Study data were excluded because the selection on abnormal glucose tolerance may impact on positive predictive value

† Categories are up to and including the latter score