

SUPPLEMENTARY DATA

Supplementary Table 1. Demographics and Clinical Characteristics of Participants

	T1D		T2D	
	N	%	N	%
Overall N	261*	100	339 ^a	100
Diabetes Duration				
<1 year	2	<1%	112	33%
1–<2 years	6	2%	74	22%
2–<3 years	101	39%	59	17%
3–<4 years	122	47%	38	11%
≥4 years	30	11%	56	17%
<i>Median (25th, 75th percentiles)</i>	3.2 (2.8, 3.7)		1.7 (0.6, 3.3)	
<i>Range</i>	0.0 – 4.7		0.0 – 8.8	
Age				
10–<13 years	91	35%	61	18%
13–<15 years	87	33%	86	25%
15–<18 years	83	32%	192	57%
<i>Mean ± SD</i>	13.9 ± 2.1		15.1 ± 2.0	
<i>Range</i>	10.0 – 17.9		10.0 – <18.0	
Gender				
Female	128	49%	219	65%
Male	133	51%	120	35%
Race/Ethnicity				
White	155	60%	26	8%
Hispanic or Latino	64	25%	194	58%
Black/African American	22	9%	94	28%
Other/Multiple race	17	7%	22	7%
Lives with Single Parent	74	28%	164	49%
Health Insurance				
Private	151	59%	89	26%
Children’s Health Plan or Other Government	87	34%	225	66%
Military	6	2%	3	<1%
None	12	5%	22	6%
Parent Education				
High School or less	89	41%	228	72%
Associate	31	14%	44	14%
Bachelor	59	27%	34	11%
Master/Professional Degree	37	17%	12	4%
Family Income				
<\$25,000	34	18%	106	46%
\$25,000–\$49,999	39	21%	73	32%
\$50,000–\$74,999	33	18%	27	12%
\$75,000–\$99,999	25	13%	13	6%
≥\$100,000	56	30%	10	4%
BMI Percentile				
<85	81	62%	10	3%
85–<95	24	18%	32	10%
95–<99	18	14%	138	43%
≥99	8	6%	142	44%
HbA1c % (mmol/mol)				
<6 (<42)	6	3%	75	24%
6–<7 (42–<53)	28	15%	65	21%

SUPPLEMENTARY DATA

7–<8 (53–<64)	58	30%	47	15%
8–<9 (64–<75)	43	22%	32	10%
≥9 (≥75)	57	30%	89	29%
<i>Mean ± SD</i>	8.3 ± 1.7		7.9 ± 2.4	
	(67 ± 18)		(63 ± 26)	
<i>Range</i>	5.5 – 15.0		4.8 – 15.5	
	(37 – 140)		(29 – 146)	
<u>Pump Use</u>	111	43%	0	0%

* Number of participants with missing or unknown data by type (Type 1/Type 2):race/ethnicity (3/3), lives with (0/3), health insurance (5/0), parent education (45/21), family income (74/110), BMI % (130/17) and HbA1c % (69/31).