

All articles in *Diabetes Care* are available on the World Wide Web at www.diabetes.org/diabetescare. Cover articles (▶) and ADA Statements are available free of charge. Other articles are available free to print subscribers or can be purchased as e-prints or reprints.

Original Research

Clinical Care/Education/Nutrition/Psychosocial Research

- 249** The DURability of Basal versus Lispro mix 75/25 insulin Efficacy (DURABLE) trial: comparing the durability of lispro mix 75/25 and glargine
J.B. Buse, B.H.R. Wolfenbittel, W.H. Herman, S. Hippler, S.A. Martin, H.H. Jiang, S.K. Shenouda, J.L. Fahrback
- 256** Randomized study of basal-bolus insulin therapy in the inpatient management of patients with type 2 diabetes undergoing general surgery (RABBIT 2 surgery)
G.E. Umpierrez, D. Smiley, S. Jacobs, L. Peng, A. Temponi, P. Mulligan, D. Umpierrez, C. Newton, D. Olson, M. Rizzo
- ▶ **262** Structured self-monitoring of blood glucose significantly reduces A1C levels in poorly controlled, noninsulin-treated type 2 diabetes: results from the Structured Testing Program study
W.H. Polonsky, L. Fisher, C.H. Schikman, D.A. Hinnen, C.G. Parkin, Z. Jelsovsky, B. Petersen, M. Schweitzer, R.S. Wagner
- 268** Better glycemic control is associated with maintenance of lower-extremity function over time in Mexican American and European American older adults with diabetes
C.-P. Wang, H.P. Hazuda
- 274** Glycemic control and health disparities in older ethnically diverse underserved adults with diabetes: five-year results from the Informatics for Diabetes Education and Telemedicine (IDEATel) study
R.S. Weinstock, J.A. Teresi, R. Goland, R. Izquierdo, W. Palmas, J.P. Eimicke, S. Ebner, S. Shea, the IDEATel Consortium
- 280** Low-dose aspirin therapy in patients with type 2 diabetes and reduced glomerular filtration rate: subanalysis from the JPAD trial
Y. Saito, T. Morimoto, H. Ogawa, M. Nakayama, S. Uemura, N. Doi, H. Jinnouchi, M. Waki, H. Soejima, S. Sugiyama, S. Okada, Y. Akai, on behalf of the Japanese Primary Prevention of Atherosclerosis with Aspirin for Diabetes (JPAD) Trial Investigators
- 286** Urinary C-peptide creatinine ratio is a practical outpatient tool for identifying hepatocyte nuclear factor 1- α /hepatocyte nuclear factor 4- α maturity-onset diabetes of the young from long-duration type 1 diabetes
R.E.J. Besser, M.H. Shepherd, T.J. McDonald, B.M. Shields, B.A. Knight, S. Ellard, A.T. Hattersley
- 292** Elevation of 1-hour plasma glucose during oral glucose tolerance testing is associated with worse pulmonary function in cystic fibrosis
J. Brodsky, S. Dougherty, R. Makani, R.C. Rubenstein, A. Kelly
- 296** Breastfeeding in women with type 1 diabetes: exploration of predictive factors
C. Sparud-Lundin, M. Wennergren, A. Elfvin, M. Berg
- 302** Effect of automated bio-behavioral feedback on the control of type 1 diabetes
B.P. Kovatchev, P. Mendosa, S. Anderson, J.S. Hawley, L.M. Ritterband, L. Gonder-Frederick

308 Improved glycemic control without hypoglycemia in elderly diabetic patients using the ubiquitous healthcare service, a new medical information system
S. Lim, S.M. Kang, H. Shin, H.J. Lee, J.W. Yoon, S.H. Yu, S.-Y. Kim, S.Y. Yoo, H.S. Jung, K.S. Park, J.O. Ryu, H.C. Jang

314 Weight-related quality of life, health utility, psychological well-being, and satisfaction with exenatide once weekly compared with sitagliptin or pioglitazone after 26 weeks of treatment
J.H. Best, R.R. Rubin, M. Peyrot, Y. Li, P. Yan, J. Malloy, L.P. Garrison

320 Web-based depression treatment for type 1 and type 2 diabetic patients: a randomized, controlled trial
K.M.P. van Bastelaar, F. Pouwer, P. Cuijpers, H. Riper, F.J. Snoek

326 Feasibility, acceptability, and predictive validity of a psychosocial screening program for children and youth newly diagnosed with type 1 diabetes
D.D. Schwartz, V.D. Cline, M.E. Axelrad, B.J. Anderson

332 Improvement in sexual functioning in patients with type 2 diabetes and depression treated with bupropion
G.S. Sayuk, B.M. Gott, B.D. Nix, P.J. Lustman

335 Suboptimal performance of blood glucose meters in an antenatal diabetes clinic
N.J. Perera, L. Molyneaux, M.I. Constantino, M. McGill, D.K.-S. Yue, S.M. Twigg, G.P. Ross

338 Illness and treatment perceptions are associated with adherence to medications, diet, and exercise in diabetic patients
E. Broadbent, L. Donkin, J.C. Stroh

Epidemiology/Health Services Research

341 Pericardial fat and echocardiographic measures of cardiac abnormalities: the Jackson Heart Study
J. Liu, C.S. Fox, D.A. Hickson, W.L. May, J. Ding, J.J. Carr, H.A. Taylor

347 Quality of diabetes care predicts the development of cardiovascular events: results of the AMD-QUASAR study
M.C.E. Rossi, G. Lucisano, M. Comaschi, C. Coscelli, D. Cucinotta, P.D. Blasi, G. Bader, F. Pellegrini, U. Valentini, G. Vespasiani, A. Nicolucci, on behalf of the AMD-QUASAR Study Group

353 Trends in the prevalence of type 2 diabetes in Asians versus whites: results from the United States National Health Interview Survey, 1997–2008
J.W.R. Lee, F.L. Brancati, H.-C. Yeh

358 Islet autoantibody seroconversion in the DPT-1 study: justification for repeat screening throughout childhood
K. Vehik, M.J. Haller, C.A. Beam, D.A. Schatz, D.K. Wherrett, J.M. Sosenko, J.P. Krischer, DPT-1 Study Group

363 Fasting plasma glucose and hemoglobin A_{1c} in identifying and predicting diabetes: the Strong Heart Study
W. Wang, E.T. Lee, B.V. Howard, R.R. Fabsitz, R.B. Devereux, T.K. Welty

369 Use of dipeptidyl peptidase-4 inhibitors and the reporting of infections: a disproportionality analysis in the World Health Organization VigiBase
M.J. Willems, A.K. Mantel-Teeuwisse, S.M. Straus, R.H. Meyboom, T.C. Egberts, H.G. Leufkens

375 Low HDL cholesterol, metformin use, and cancer risk in type 2 diabetes: the Hong Kong Diabetes Registry
X. Yang, W.Y. So, R.C.W. Ma, A.P.S. Kong, H.M. Lee, L.W.L. Yu, C.-C. Chow, R. Ozaki, G.T.C. Ko, J.C.N. Chan

381 Association between periodontitis and impaired fasting glucose and diabetes
Y.-H. Choi, R.E. McKeown, E.J. Mayer-Davis, A.D. Liese, K.-B. Song, A.T. Merchant

Table of Contents continues on page vii

387 Implications of alternative definitions of prediabetes for prevalence in U.S. adults

C. James, K.M. Bullard, D.B. Rolka, L.S. Geiss, D.E. Williams, C.C. Cowie, A. Albright, E.W. Gregg

392 Endotoxemia is associated with an increased risk of incident diabetes

P.J. Pussinen, A.S. Havulinna, M. Lehto, J. Sundvall, V. Salomaa

398 Risk of acute complications of diabetes among people with schizophrenia in Ontario, Canada

T. Becker, J. Hux

403 HbA_{1c} in nondiabetic Dutch infants aged 8–12 months: the GECKO-Drenthe birth cohort study

H. Jansen, H.G. Huiting, S. Scholtens, P.J.J. Sauer, R.P. Stolk

Emerging Treatments and Technologies

406 Closed-loop insulin delivery during pregnancy complicated by type 1 diabetes

H.R. Murphy, D. Elleri, J.M. Allen, J. Harris, D. Simmons, G. Rayman, R. Temple, D.B. Dunger, A. Haidar, M. Nodale, M.E. Wilinska, R. Hovorka

Pathophysiology/Complications

► **412** Glucagon-like peptide-1 receptor agonist treatment prevents glucocorticoid-induced glucose intolerance and islet-cell dysfunction in humans

D.H. van Raalte, R.E. van Genugten, M.M.L. Linssen, D.M. Ouwens, M. Diamant

418 Impact of positive family history and genetic risk variants on the incidence of diabetes: the Finnish Diabetes Prevention Study

M.I. Uusitupa, A. Stančáková, M. Peltonen, J.G. Eriksson, J. Lindström, S. Aunola, P. Ilanne-Parikka, S. Keinänen-Kiukaanniemi, J. Tuomilehto, M. Laakso

424 High rate of regression from micro-macroalbuminuria to normoalbuminuria in children and adolescents with type 1 diabetes treated or not with enalapril: the influence of HDL cholesterol

S. Salardi, C. Balsamo, S. Zucchini, G. Maltoni, M. Scipione, A. Rollo, S. Gualandi, A. Cicognani

430 Body and liver fat mass rather than muscle mitochondrial function determine glucose metabolism in women with a history of gestational diabetes mellitus

T. Prikoszovich, C. Winzer, A.I. Schmid, J. Szendroedi, M. Chmelik, G. Pacini, M. Krššák, E. Moser, T. Funahashi, W. Waldhäusl, A. Kautzky-Willer, M. Roden

437 Insulin infusion during normoglycemia modulates insulin secretion according to whole-body insulin sensitivity

C. Anderwald, A. Tura, A. Grassi, M. Krebs, J. Szendroedi, M. Roden, M.G. Bischof, A. Luger, G. Pacini

► **442** Higher plasma levels of advanced glycation end products are associated with incident cardiovascular disease and all-cause mortality in type 1 diabetes: a 12-year follow-up study

J.W. Nin, A. Jorsal, I. Ferreira, C.G. Schalkwijk, M.H. Prins, H.-H. Parving, L. Tarnow, P. Rossing, C.D. Stehouwer

448 Liver ATP synthesis is lower and relates to insulin sensitivity in patients with type 2 diabetes

A.I. Schmid, J. Szendroedi, M. Chmelik, M. Krššák, E. Moser, M. Roden

454 Vitamin D deficiency and coronary artery calcification in subjects with type 1 diabetes

K.A. Young, J.K. Snell-Bergeon, R.G. Naik, J.E. Hokanson, D. Tarullo, P.A. Gottlieb, S.K. Garg, M. Rewers

459 Cerebral perfusion and aortic stiffness are independent predictors of white matter brain atrophy in type 1 diabetic patients assessed with magnetic resonance imaging

S.G.C. van Elderen, A. Brandts, J. van der Grond, J.J.M. Westenberg, L.J.M. Kroft, M.A. van Buchem, J.W.A. Smit, A. de Roos

464 The prevalence and predictors of an abnormal ankle-brachial index in the Bypass Angioplasty Revascularization Investigation 2 Diabetes (BARI 2D) Trial

P.P. Singh, J.D. Abbott, M.S. Lombardero, K. Sutton-Tyrrell, G. Woodhead, L. Venkitachalam, N.P. Tsapatsaris, T.C. Piemonte, R.M. Lago, M.K. Rutter, R.W. Nesto, BARI 2D Study Group

468 Relationship between carotid intima-media thickness and the presence and extent of coronary stenosis in type 2 diabetic patients with carotid atherosclerosis but without history of coronary artery disease

R. Kasami, H. Kaneto, N. Katakami, S. Sumitsuji, K. Yamasaki, T. Kuroda, K. Tachibana, T. Yasuda, A. Kuroda, T.-a. Matsuoka, M. Matsuhisa, I. Shimomura

471 Clinical and genetic characteristics of patients with type 1 diabetes associated with interferon therapy

K. Nakanishi, S. Saitoh

474 Serum apolipoprotein AI and B are stronger biomarkers of diabetic retinopathy than traditional lipids

M.B. Sasongko, T.Y. Wong, T.T. Nguyen, R. Kawasaki, A. Jenkins, J. Shaw, J.J. Wang

Cardiovascular and Metabolic Risk

480 The association between maternal glucose concentration and child BMI at age 3 years

A.L. Deierlein, A.M. Siega-Riz, K. Chantala, A.H. Herring

485 Ethnic differences in pancreatic fat accumulation and its relationship with other fat depots and inflammatory markers

K.-A. Lê, E.E. Ventura, J.Q. Fisher, J.N. Davis, M.J. Weigensberg, M. Punyanitya, H.H. Hu, K.S. Nayak, M.I. Goran

491 Urbanization, physical activity, and metabolic health in sub-Saharan Africa

F.K. Assah, U. Ekelund, S. Brage, J.C. Mbanya, N.J. Wareham

497 Sedentary activity associated with metabolic syndrome independent of physical activity

A. Bankoski, T.B. Harris, J.J. McClain, R.J. Brychta, P. Caserotti, K.Y. Chen, D. Berrigan, R.P. Troiano, A. Koster

504 Distribution of abdominal visceral and subcutaneous adipose tissue and metabolic syndrome in a Korean population

S. Kim, B. Cho, H. Lee, K. Choi, S.S. Hwang, D. Kim, K. Kim, H. Kwon

507 The relation of diabetes, impaired fasting blood glucose, and insulin resistance to left ventricular structure and function in African Americans: the Jackson Heart Study

E.R. Fox, D.F. Sarpong, J.C. Cook, T.E. Samdarshi, H.S. Nagarajarao, P.R. Liebson, M. Sims, G. Howard, R. Garrison, H.A. Taylor

Reviews/Commentaries/ADA Statements

510 Meta-analysis

Efficacy of insulin analogs in achieving the hemoglobin A_{1c} target of <7% in type 2 diabetes: meta-analysis of randomized controlled trials

D. Giugliano, M.I. Maiorino, G. Bellastella, P. Chiodini, A. Ceriello, K. Esposito

518 Review

A1C versus glucose testing: a comparison

D.B. Sacks

Editorials

524 Editorial (see Sacks, p. 518)

Diagnosing diabetes with glucose criteria: worshipping a false god

M.B. Davidson

527 Editorial (see Polonsky et al., p. 262)

Home blood glucose monitoring in type 2 diabetes: broken health care system undermines study's impact
I.B. Hirsch

529 Editorial (see Sasongko et al., p. 474)

Apolipoproteins and diabetic retinopathy
A. Chait, V.N. Montes

532 Issues and Events

Diabetes Care Electronic Pages

Available at <http://care.diabetesjournals.org>

Perspectives on the News

e11 Type 2 diabetes: uses of thiazolidinediones and insulin
Z.T. Bloomgarden

Online Letters: Observations

e17 Glycemic response to ingested Dreamfields pasta compared with traditional pasta
F.Q. Nuttall, M.C. Gannon, H. Hoover

e19 HDL dysfunctionality (paraoxonase) is worse in nondiabetic, postmenopausal African American than in white women
T. Gaillard, S. Parthasarathy, K. Osei

Online Letters: Comments and Responses

e20 Comment on: Zhang et al. A1C level and future risk of diabetes: a systematic review. *Diabetes Care* 2010;33:1665–1673
C.A. Robinson, M. Abdel-Motagally

e21 Response to comment on: Zhang et al. A1C level and future risk of diabetes: a systematic review. *Diabetes Care* 2010;33:1665–1673

X. Zhang, E.W. Gregg, D.F. Williamson, L.E. Barker, W. Thomas, K.M. Bullard, G. Imperatore, D.E. Williams, A.L. Albright

e22 Comment on: Luijf et al. Premeal injection of rapid-acting insulin reduces postprandial glycemic excursions in type 1 diabetes. *Diabetes Care* 2010;33:2152–2155

A.E. Scaramuzza, D. Iafusco, A. Bosetti, A.D. Palma, G.V. Zuccotti

e23 Response to comment on: Luijf et al. Premeal injection of rapid-acting insulin reduces postprandial glycemic excursions in type 1 diabetes. *Diabetes Care* 2010;33:2152–2155

Y.M. Luijf, J.H. DeVries