

SUPPLEMENTARY DATA

Supplementary Table 1. Incidence rate of diagnosed diabetes by quartiles of NT-proBNP at baseline.

NT-proBNP quartile [Range, pg/mL] N	Q1 [<31] N=1953	Q2 [31.1-61.8] N=1952	Q3 [61.9-114.4] N=1957	Q4 [>114.5] N=1960
Person-years of follow up	19489.19	19804.26	19956.01	19848.89
Cases of Diagnosed Diabetes	539	461	397	343
Incidence rate (per 1000 person-years)	27.66	23.28	19.89	17.28

Supplementary Table 2. Adjusted^{fl} Hazard Ratios (95% Confidence Intervals) of Incident Diagnosed Diabetes by Quartile of NT-proBNP at Baseline After Additional Exclusions.

NT-proBNP quartile [Range, pg/mL]	Q1 [<31]	Q2 [31.1-61.8]	Q3 [61.9-114.4]	Q4 [>114.5]
Excluding people with elevated hs-cTnt N=386	1.0 Reference	0.84 (0.73-0.96)	0.78 (0.68-0.90)	0.71 (0.61-0.83)
Excluding people with undiagnosed diabetes N=438	1.0 Reference	0.88 (0.76-1.02)	0.80 (0.69-0.94)	0.71 (0.60-0.84)

^a Adjusted for: Age, sex, race, ARIC center, education, BMI, height², SBP, DBP, smoking, LDL, HDL, hs-CRP, fasting glucose and family history of diabetes