


# diabetes

A JOURNAL OF THE  
AMERICAN DIABETES  
ASSOCIATION®

[www.diabetes.org/diabetes](http://www.diabetes.org/diabetes)



## SIGNAL TRANSDUCTION

Ceramide- and oxidant-induced insulin resistance involve loss of insulin-dependent Rac-activation and actin remodeling in muscle cells

L. JEBAILY, O. WANONO, W. NIU, J. ROESSLER, A. RUDICH, AND A. KLIP 394

Protein-tyrosine phosphatase 1B-deficient myocytes show increased insulin sensitivity and protection against tumor necrosis factor- $\alpha$ -induced insulin resistance

I. NIETO-VAZQUEZ, S. FERNÁNDEZ-VELEDO, C. DE ALVARO, C.M. RONDINONE, A.M. VALVERDE, AND M. LORENZO 404

The Rab GTPase-activating protein AS160 integrates Akt, protein kinase C, and AMP-activated protein kinase signals regulating GLUT4 traffic

F.S.L. THONG, P.J. BILAN, AND A. KLIP 414

## OBESITY STUDIES

IGF-binding protein-2 protects against the development of obesity and insulin resistance

S.B. WHEATCROFT, M.T. KEARNEY, A.M. SHAH, V.A. EZZAT, J.R. MIELL, M. MODO, S.C.R. WILLIAMS, W.P. CAWTHORN, G. MEDINA-GOMEZ, A. VIDAL-PUIG, J.K. SETHI, AND P.A. CROSSEY 285

Wnt10b inhibits obesity in *ob/ob* and agouti mice

W.S. WRIGHT, K.A. LONGO, V.W. DOLINSKY, I. GERIN, S. KANG, C.N. BENNETT, S.-H. CHIANG, T.C. PRESTWICH, C. GRESS, C.F. BURANT, V.S. SUSULIC, AND O.A. MACDOUGALD 295

## IMMUNOLOGY AND TRANSPLANTATION

Cellular expression requirements for inhibition of type 1 diabetes by a dominantly protective major histocompatibility complex haplotype

Y.-G. CHEN, P.A. SILVEIRA, M.A. OSBORNE, H.D. CHAPMAN, AND D.V. SERREZ 424

## ISLET STUDIES

Identification of the amyloid-degrading enzyme neprilysin in mouse islets and potential role in islet amyloidogenesis

S. ZRAIKA, R.L. HULL, J. UDAYASANKAR, A. CLARK, K.M. UTZSCHNEIDER, J. TONG, F. GERCHMAN, AND S.E. KAHN 304

Melanin concentrating hormone is a novel regulator of islet function and growth

P. PISSIOS, U. OZCAN, E. KOKKOTOU, T. OKADA, C.W. LIEW, S. LIU, J.N. PETERS, G. DAHLGREN, J. KARAMCHANDANI, Y.C. KUDVA, A.J. KURPAD, R.T. KENNEDY, E. MARATOS-FLIER, AND R.N. KULKARNI 311

Glucose-dependent regulation of  $\gamma$ -aminobutyric acid ( $GABA_A$ ) receptor expression in mouse pancreatic islet  $\alpha$ -cells

S.J. BAILEY, M.A. RAVIER, AND G.A. RUTTER 320

An ATP-binding mutation (G334D) in *KCNJ11* is associated with a sulfonylurea-insensitive form of developmental delay, epilepsy, and neonatal diabetes

R. MASIA, J.C. KOSTER, S. TUMINI, F. CHIARELLI, C. COLOMBO, C.G. NICHOLS, AND F. BARBETTI 328

Glucotoxicity in the INS-1 rat insulinoma cell line is mediated by the orphan nuclear receptor small heterodimer partner

K.-G. PARK, K.-M. LEE, H.-Y. SEO, J.-H. SUH, H.-S. KIM, L. WANG, K.-C. WON, H.-W. LEE, J.-Y. PARK, K.-U. LEE, J.-G. KIM, B.-W. KIM, H.-S. CHOI, AND I.-K. LEE 431

Sulfonylurea compounds uncouple the glucose dependence of the insulinotropic effect of glucagon-like peptide 1

J. DE HEER AND J.J. HOLST 438

## PATHOPHYSIOLOGY

Pancreatic lipomatosis is a structural marker in nondiabetic children with mutations in carboxyl-ester lipase

H. RÆDER, IS. HALDRSEN, L. ERSLAND, R. GRÜNER, T. TAXT, O. SØVIK, A. MOLVEN, AND P.R. NJØLSTAD 444

Tumor necrosis factor- $\alpha$  induces intestinal insulin resistance and stimulates the overproduction of intestinal apolipoprotein B48-containing lipoproteins

B. QIN, W. QIU, R.K. AVRAMOGLU, AND K. ADELI 450

## COMPLICATIONS

Salicylate-based anti-inflammatory drugs inhibit the early lesion of diabetic retinopathy

L. ZHENG, S.J. HOWELL, D.A. HATALA, K. HUANG, AND T.S. KERN 337

Deletion of protein kinase C- $\beta$  isoform in vivo reduces renal hypertrophy but not albuminuria in the streptozotocin-induced diabetic mouse model

M. MEIER, J.-K. PARK, D. OVERHEU, T. KIRSCH, C. LINDSCHAU, F. GUELER, M. LEITGES, J. MENNE, AND H. HALLER 346

Treatment of streptozotocin-induced diabetic rats with AVE7688, a vasopeptidase inhibitor: effect on vascular and neural disease

E.P. DAVIDSON, T.L. KLEINSCHMIDT, C.L. OLTMAN, D.D. LUND, AND M.A. YOREK 355

Macrophage scavenger receptor-A-deficient mice are resistant against diabetic nephropathy through amelioration of microinflammation

H.K. USUI, K. SHIKATA, M. SASAKI, S. OKADA, M. MATSUDA, Y. SHIKATA, D. OGAWA, Y. KIDO, R. NAGASE, K. YOZAI, S. OHGA, A. TONE, J. WADA, M. TAKEYA, S. HORIUCHI, T. KODAMA, AND H. MAKINO 363

Topical administration of Nepafenac inhibits diabetes-induced retinal microvascular disease and underlying abnormalities of retinal metabolism and physiology T.S. KERN, C.M. MILLER, Y. DU, L. ZHENG, S. MOHR, S.L. BALL, J.A. JAMISON, M. KIM, AND D.P. BINGAMAN	373	P2 promoter variants of the hepatocyte nuclear factor 4 $\alpha$ gene are associated with type 2 diabetes in Mexican Americans D.M. LEHMAN, D.K. RICHARDSON, C.P. JENKINSON, K.J. HUNT, T.D. DYER, R.J. LEACH, R. ARYA, H.E. ABBoud, J. BLANGERO, R. DUGGIRALA, AND M.P. STERN	513
Evidence for a role of transforming growth factor (TGF)- $\beta$ 1 in the induction of postglomerular albuminuria in diabetic nephropathy: amelioration by soluble TGF- $\beta$ type II receptor L.M. RUSSO, E. DEL RE, D. BROWN, AND H.Y. LIN	380	Women with mitochondrial haplogroup N9a are protected against metabolic syndrome M. TANAKA, N. FUKU, Y. NISHIGAKI, H. MATSUO, T. SEGAWA, S. WATANABE, K. KATO, K. YOKO, M. ITO, Y. NOZAWA, AND Y. YAMADA	518
Alterations of adenine nucleotide metabolism and function of blood platelets in patients with diabetes A. MICHNO, H. BIELARCZYK, T. PAWEŁCZYK, A. JANKOWSKA-KULAWY, J. KLIMASZEWSKA, AND A. SZUTOWICZ	462	<i>PTPN22</i> R620W functional variant in type 1 diabetes and autoimmunity related traits C. CHELALA, S. DUCHATELET, M.-L. JOFFRET, R. BERGHOLDT, D. DUBOIS-LAFORGUE, P. GHANDIL, F. POCLOT, S. CAILLAT-ZUCMAN, J. TIMSIT, AND C. JULIER	522
Impact of type 2 diabetes on nitric oxide and adrenergic modulation of myocardial perfusion M.M. MADSEN, H.M. SØNDERGAARD, M. BØTTCHER, N.H. BUUS, O. SCHMITZ, H.E. BØTKER, AND T.T. NIELSEN	468	Myocardial uptake of circulating triglycerides in nondiabetic patients with heart disease R.H. NELSON, A. PRASAD, A. LERMAN, AND J.M. MILES	527
High glucose, high insulin, and their combination rapidly induce laminin- $\beta$ 1 synthesis by regulation of mRNA translation in renal epithelial cells M.M. MARIAPPAN, D. FELIERS, S. MUMMIDI, G.G. CHOUDHURY, AND B.S. KASINATH	476	Type 2 diabetes-associated missense polymorphisms <i>KCNJ11</i> E23K and <i>ABCC8</i> A1369S influence progression to diabetes and response to interventions in the Diabetes Prevention Program J.C. FLOREZ, K.A. JABLONSKI, S.E. KAHN, P.W. FRANKS, D. DABELEA, R.F. HAMMAN, W.C. KNOWLER, D.M. NATHAN, AND D. ALTSHULER FOR THE DIABETES PREVENTION PROGRAM RESEARCH GROUP	531
<b>PHARMACOLOGY AND THERAPEUTICS</b>			
Oral insulin-mimetic compounds that act independently of insulin S. GARCÍA-VICENTE, F. YRAOLA, L. MARTÍ, E. GONZÁLEZ-MUÑOZ, M.J. GARCÍA-BARRALDO, C. CANTÓ, A. ABELLA, S. BOUR, R. ARTUCH, C. SIERRA, N. BRANDI, C. CARPÉNÉ, J. MORATINOS, M. CAMPS, M. PALACÍN, X. TESTAR, A. GUMÀ, F. ALBERICIO, M. ROYO, A. MIAN, AND A. ZORZANO	486	Linkage and association analyses of type 2 diabetes/impaired glucose metabolism and adiponectin serum levels in Japanese Americans from Hawaii I.P. KOVAC, R.J. HAVLIK, D. FOLEY, R. PEILA, D. HERNANDEZ, F. WAVRANT-DE VRIÈZE, A. SINGLETON, J. EGAN, D. TAUB, B. RODRIGUEZ, K. MASAKI, J.D. CURB, W.Y. FUJIMOTO, AND A.F. WILSON	537
<b>GENETICS</b>			
A possible role for the <i>PPARG</i> Pro12Ala polymorphism in preterm birth A. MEIRHAEGHE, C.A.G. BOREHAM, L.J. MURRAY, F. RICHARD, G.D. SMITH, I.S. YOUNG, AND P. AMOUYEL	494	Diabetes and suppressors of cytokine signaling proteins S.G. RØNN, N. BILLESTRUP, AND T. MANDRUP-POULSEN	541
Tag polymorphisms at the <i>A20</i> ( <i>TNFAIP3</i> ) locus are associated with lower gene expression and increased risk of coronary artery disease in type 2 diabetes W. BOONYASRISAWAT, D. EBERLE, S. BACCI, Y.-Y. ZHANG, D.N. NOLAN, E.V. GERVINO, M.T. JOHNSTONE, V. TRISCHITTA, S.E. SHOELSON, AND A. DORIA	499	<b>ERRATA</b>	
Genetic reconstitution of autoimmune type 1 diabetes with two major susceptibility genes in the rat N. YOKOI, C. HAYASHI, Y. FUJIWARA, H.-Y. WANG, AND S. SEINO	506	<b>ORGANIZATION SECTION</b>	
<b>SI UNITS TABLE</b>			
<b>BRIEF REPORTS</b>			
Haplotypes of transcription factor 7-like 2 ( <i>TCF7L2</i> ) gene and its upstream region are associated with type 2 diabetes and age of onset in Mexican Americans D.M. LEHMAN, K.J. HUNT, R.J. LEACH, J. HAMILTON, R. ARYA, H.E. ABBoud, R. DUGGIRALA, J. BLANGERO, H.H.H. GÖRING, AND M.P. STERN	389		